

Leucocyte Typing V

White Cell Differentiation Antigens

**Proceedings of the Fifth
International Workshop
and Conference
Held in Boston, USA
3-7 November, 1993**

Volume One

Edited by

**Stuart F. Schlossman, Laurence Boumsell, Wally Gilks,
John M. Harlan, Tadamitsu Kishimoto, Chikao Morimoto,
Jerome Ritz, Stephen Shaw, Roy Silverstein, Timothy Springer,
Thomas F. Tedder and Robert F. Todd**

**Oxford New York Tokyo
OXFORD UNIVERSITY PRESS
1995**

BEST AVAILABLE COPY

Oxford University Press, Walton Street, Oxford OX1 2DP

Oxford New York
Athens Auckland Bangkok Bombay
Calcutta Cape Town Dar es Salaam Delhi
Florence Hong Kong Istanbul Karachi
Kuala Lumpur Madras Madrid Melbourne
Mexico City Nairobi Paris Singapore
Taipei Tokyo Toronto
and associated companies in
Berlin Ibadan

Oxford is a trade mark of Oxford University Press

Published in the United States
by Oxford University Press Inc., New York

© The Organizing Committee of the Fifth International Conference on
Human Leucocyte Differentiation Antigens and Oxford University Press, 1995

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, without the prior permission in writing of Oxford University Press. Within the UK, exceptions are allowed in respect of any fair dealing for the purpose of research or private study, or criticism or review, as permitted under the Copyright, Designs and Patents Act, 1988, or in the case of reprographic reproduction in accordance with the terms of licences issued by the Copyright Licensing Agency. Enquiries concerning reproduction outside these terms and in other countries should be sent to the Rights Department, Oxford University Press, at the address above.

This book is sold subject to the condition that it shall not, by way of trade or otherwise, be lent, re-sold, hired out, or otherwise circulated without the publisher's prior consent in any form of binding or cover other than that in which it is published and without a similar condition including this condition being imposed on the subsequent purchaser.

A catalogue record for this book is available from the British Library

Library of Congress Cataloging in Publication Data
(Data available)

ISBN 0 19 2626384 Volume 1
ISBN 0 19 2623761 Two volume set
(Available only as a two volume set)

Typeset by Dobbie Typesetting Limited, Tavistock, Devon
Printed in Great Britain by
Butler and Tanner Ltd, Frome, Somerset

BEST AVAILABLE COPY

identical pattern of tyrosine phosphorylated polypeptides was observed. Prominent phosphorylated polypeptides were identified with M_r of 110, 72, and 40 kDa. Incubation of K-562 cells with MA73 (2ZC115) failed to induce a similar pattern of phosphorylated polypeptides. Incubation with the cross-linking antibody alone, GAM F(ab')₂, similarly did not induce novel phosphorylated polypeptides.

The pattern of tyrosine phosphorylation observed following cross-linking of anti-CD32 mAb is consistent with data previously reported utilizing Fab fragments of mAb IV.3 that showed that the 40-kDa tyrosine phosphorylated protein was FcγRII [8]. The identities of the other tyrosine phosphorylated substrates of M_r 110 and 72 kDa are currently under investigation.

The observation that mAb MA23 (BAS62-11) induced a similar pattern of tyrosine phosphorylated polypeptides but yet does not recognize CD32 suggests that this mAb binds an antigen on the surface of K-562 cells via the Fab domain and then activates FcγRII via its Fc region. This could be accomplished by: (1) formation of cellular immune complexes that could bind to FcγRII on other K-562 cells; or (2) tripartite engagement of IgG molecules on the same cell with subsequent cross-linking by the secondary antibody. This finding indicates that ascites containing whole immunoglobulin of an mAb directed against a different cell surface molecule could induce FcγRII-mediated tyrosine phosphorylation. Thus it points out the necessity of using Fab or F(ab')₂ fragments of mAb when investigating the cellular signal

transduction mechanisms of any receptors on cells expressing FcγR. The physical cross-linking of such intact immunoglobulin molecules may produce patterns of tyrosine phosphorylated proteins similar to those induced by cross-linking of FcγRII alone.

Acknowledgement

These studies were supported by NIH grant CA38055.

References

1. Ravetch, J. V. and Klein, J.-P. *Ann. Rev. Immunol.* 9, 457 (1991).
2. Brooks, D. G., Qui, W. Q., Luster, A. L., and Ravetch, J. V. *J. exp. Med.* 170, 1369 (1989).
3. Huang, M., Indik, Z., Brass, L. F., Hoxie J. A., Schreiber, A. D., and Brugge, J. S. *J. biol. Chem.* 267, 3467 (1992).
4. Scholl, P. R., Ahern, D., and Geha, R. S. *J. Immunol.* 149, 1751 (1992).
5. Liao, F., Shin, H. S., and Rhee, S. G. *Proc. natl Acad. Sci., USA* 89, 3659 (1992).
6. Sakudo, T. W. and Flett, H. B. *Cell Prolif.* 24, 383 (1991).
7. Looney, R. J., Abraham, G. N., and Anderson, C. L. *J. Immunol.* 136, 1641 (1986).
8. Gharizadeh, S. and Flett, H. B. *J. Immunol.* 153, 30 (1994).
9. Flett, H. B. *Clin. Immunol. Immunopathol.* 59, 222 (1991).

M8.2 Specificity of CD32 mAb for FcγRIIa, FcγRIIb1, and FcγRIIb2 expressed in transfected mouse B cells and BHK-21 cells

PETRA BUDDE, VOLKER WEINRICH, PETER SONDERMANN, NILS BEWARDER, ANDREAS KILIAN, OLAF SCHULZECK, and JÜRGEN FREY

Six monoclonal antibodies (mAb) of the CD32 panel were analysed for their specificity against the various FcγRII isoforms expressed in the FcγR - mouse B-cell line IIA1.6 [1] and BHK-21 cells [2]. In addition, we compared the reactivity of the mAb with the respective receptors homologously expressed in the human B-cell line Daudi (FcγRIIb1 + and FcγRIIb2 +) as well as K-562 cells (FcγRIIaHR + /LR +; HR = high

responder and LR = low responder). Besides the six Workshop antibodies, we included three new mAb obtained in our laboratory, 1A4, IIA5, and IIBD2, which were compared with an mAb MA179 (AT10) known to recognize all CD32 isoforms [3].

Using FACS analysis, we found that the two mAb MA23 (BAS62-11) and MA73 (2ZC115) did not react with any of the FcγRII, independent of the cell lines

⚠ NOTICE THIS MATERIAL MAY BE PROTECTED BY COPYRIGHT LAW (TITLE 17 U.S. CODE)

ATTORNEY DOCKET NUMBER: 11183-003-999
SERIAL NUMBER: 10/524,134
REFERENCE: C09

BEST AVAILABLE COPY

Table 1 Reactivity of mAb with homologically and heterologously expressed CD32 isotypes

CD32 isoforms ¹	Reactivity of mAb ²									
	MA179 (AT10)	MR7 (IV.3)	MA23 (BAS62-11)	MA72 (KB61)	MA73 (Z2C115)	MA128 (PILR.26)	MA126 (CDM45)	IA4	II1A5	II8D2
IIA1.6 cell lines										
FcγRIIaLR	++	++	0	++	0	+	++	0	0	0
FcγRIIaHR	+++	++	0	++	0	+	+	0	0	0
FcγRIIb1	++	0	0	+++	0	+	0	0	0	0
FcγRIIb2	++	0	0	+++	0	+	0	0	0	0
BHK-21 cell lines										
FcγRIIaLR	++	++	0	+++	0	++	++	0	++	+
FcγRIIaHR	+++	++	0	+++	0	++	++	0	+	ND
FcγRIIb1	+++	++	0	+++	0	++	++	0	++	++
FcγRIIb2	+++	++	0	+++	0	++	++	0	++	++
Other cell lines										
Daudi	+++	(±)	0	+++	0	++	+	++	0	0
K-562	+++	+++	0	+++	0	++	+++	0	0	0

¹The FcγRII isoforms were transfected into FcγR - mouse B-cell line II1A1.6 and BHK-21 cells as described [1]. mAb were incubated with stable clones expressing the respective isoforms under saturating conditions.

²Bound mAb were detected using fluorescein isothiocyanate (FITC)-labelled F(ab')₂ fragment of goat anti-mouse IgG + IgM. Background fluorescence was detected using the respective mouse isotype controls (IgG/IgM) and the Fcγ-paramal cell lines. 0, No reactivity; ±, very weak reactivity; + to +++ indicate increasing levels of reactivity; ND, not done.

BEST AVAILABLE COPY




Fig. 1 Reactivity of CD32 mAb on CD19+ human B cells prepared from tonsils. After T-cell rosetting 98 per cent of the cells were CD19+. Cells (8×10^5) were incubated with the various Workshop mAb (1:100 diluted) and mAb AT10 and 1A4 (Fc γ RIIb-specific) as culture supernatants, followed by incubation with a fluorescein isothiocyanate (FITC)-labelled goat anti-mouse IgG+IgM F(ab')₂ fragment and analysed by flow cytometry.

studied (Table 1). mAb MA128 (FL18.26) recognizes Fc γ RIIa, Fc γ RIIb1, and Fc γ RIIb2 equally well, independently of the cell line studied. In contrast, mAb MA72 (KB61) shows a preferential binding to Fc γ RIIb1 and Fc γ RIIb2 in IIA1.6 cells (Table 1). mAb MR7 (IV.3) and MA126 (CIKM5) showed a strong preferential binding to Fc γ RIIa compared to Fc γ RIIb1 and Fc γ RIIb2, when expressed either in mouse or human B cells. Interestingly, this could only be observed in mouse B cells (IIA1.6) but not in BHK-21 cells. In this cell line the Fc γ RIIb isoforms are also recognized by MR7 (IV.3) and MA126 (CIKM5). Therefore, either different glycosylation patterns of the respective Fc γ RIIb isoforms or associated surface molecules in B cells are responsible for the varying antibody specificity. None of the mAb reacted with CD16-Fc γ RIIa chimeric receptors containing either 23 or 47 amino acids (aa) of the extracellular region of Fc γ RIIa (plus transmembrane and cytoplasmic region) [2]. Among the Workshop antibodies tested on human tonsillar B cells, only mAb MA72 (KB61) and MA128 (FL18.26) gave positive results (Fig. 1). In contrast to all transfected cell lines analysed as well as Daudi and K-562 cells, MA23 (BAS62-11) gave bright fluorescence signals on human tonsillar B cells (Fig. 1).

Using a synthetic peptide (aa 30-39 of the mature protein) of Fc γ RIIb2 as well as Fc γ RIIb2 expressed in *Escherichia coli* we raised a panel of mAb with varying specificity. mAb 1A4 (IgM) directed against the synthetic peptide shows a strong specificity for Fc γ RIIb expressed in human B cells and B-cell lines comparable to that of mAb MA179 (AT10) and MA72 (KB61) (Fig. 1; Table 1). Interestingly, this mAb does not react with Fc γ RIIb1 and Fc γ RIIb2 expressed in mouse B cells (IIA1.6) as well as in BHK-21 cells (Table 1). Further studies (not described) revealed, that mAb 1A4 mostly reacts with activated B cells. The specificity of the antibody was verified by immunoprecipitation of Fc γ RIIb1 and Fc γ RIIb2 from Daudi cells (Table 2). The mAb II1A5 and II8D2 were raised against the Fc γ RIIb2 expressed in *E. coli* and were selected on BHK-21 cells expressing Fc γ RIIb2. In FACS analyses mAb II1A5 and II8D2 recognize Fc γ RIIa and Fc γ RIIb isoforms only when they are expressed on BHK-21 cells. In contrast, in Western blot analyses both antibodies detected Fc γ RII, independently of the cell line expressing the receptors (Table 2). Here, mAb II8D2 shows specificity for the Fc γ RIIb isoforms, whereas mAb II1A5 recognizes both Fc γ RIIa and Fc γ RIIb. Thus, it is possible to differentiate between Fc γ RIIa and Fc γ RIIb isoforms expressed in different cells and cell lines by Western blot analysis.




Fig. 2 Immunoprecipitation of homologously and heterologously expressed CD32 using mAb FL18.26. Immunoprecipitation and detection were performed as described in the legend to Table 2. The figure shows a representative result of the precipitation experiments summarized in Table 2.

The efficiency of the Workshop antibodies for immunoprecipitation of Fc γ RII isoforms from different cells was analysed using transfected BHK-21 and IIA1.6 cells as well as Daudi cells. The Fc γ RII precipitation was judged by immunoblotting using the new mAb IIAS. Among the antibodies tested, only mAb AT10 and FL18.26 were able to bind both Fc γ RIIa and Fc γ RIIb isoforms with affinities sufficient to isolate the immune complexes (Table 2). These results confirm the data obtained by FACS analysis (Table 1). Using mAb MR7 (IV.3) we could only isolate the Fc γ RIIa from BHK-21 and IIA1.6 cells. The reactivity of MR7 (IV.3) against the Fc γ RIIb isoforms expressed in BHK-21 cells observed by FACS analysis (Table 1) must be a fairly weak binding because we could not isolate these Fc γ RII by immunoprecipitation (Table 2). Comparable results were obtained with mAb CIKMS. The only difference is that CIKMS is more efficient in immunoprecipitating the Fc γ RIIaLR alloform (Table 2). The counterpart to MR7 (IV.3) and MA126 (CIKMS) for immunoprecipitation is mAb MA72 (KB61), which specifically reacts with the Fc γ RIIb isoforms (Table 2). This differential reactivity is not

Table 2 Immunoprecipitation efficiency of anti-CD32 mAb with homologously and heterologously expressed receptor isoforms*

mAb		Immunoprecipitation efficiency with [†]						
Workshop code	Clone name	BHK-21		IIA1.6		BHK-21	IIA1.6	Daudi
		Fc γ RIIaHR	Fc γ RIIaLR	Fc γ RIIaHR	Fc γ RIIaLR	Fc γ RIIb2	Fc γ RIIb2	Fc γ RIIb
MR7	IV.3	+++	+++	+++	+++	-	-	-
MA23	BAS62-11	-	-	-	-	-	-	-
MA72	KB61	±	±	±	±	+++	+++	+
MA73	2ZC115	-	-	-	-	+++	+++	+
MA128	FL18.26	++	++	++	++	++	++	+
MA126	CIKMS	++	+++	+	+++	-	-	-
MA179	AT10	+++	+++	+++	+++	++	++	+
	IIIA5	++	++	-	-	++	-	-
	IIID2	++	-	-	-	++	-	-
	IA4	ND	ND	ND	ND	ND	ND	ND

*Cells (see footnote to Table 1) were incubated with the mAb under saturating conditions at 4 °C (except for mAb MA73 where the cells were lysed before adding the antibody). The cells were subsequently lysed in modified RIPA buffer (10 mM Tris-HCl, pH 7.2; 1% w/v Triton-X-100; 1% sodium deoxycholate; 0.1% sodium dodecyl sulfate (SDS); 150 mM NaCl; 5 mM Na-EDTA; 4 mM phenylmethylsulfonyl fluoride (PMSF); 1 TIU/ml aprotinin). The cell-free supernatant was subjected to Protein A + Q-Sepharose (90 min, 4 °C). Bound immune complexes were eluted using sample buffer and were subjected to SDS-polyacrylamide gel electrophoresis (PAGE). After blotting on to nitrocellulose membranes, the various Fc γ RII isoforms were detected using mAb IIID2 (Fc γ RIIb) and IIIA5 (Fc γ RIIa + Fc γ RIIb). Bound mAb was detected after incubation with peroxidase-labelled goat anti-mouse IgG + IgM using the ECL chemiluminescence detection system (Amersham).

†ND, Not done. -, No reaction; ±, very weak reactivity; + to +++ indicates increasing levels of reactivity.

BEST AVAILABLE COPY

observed in FACS analyses, which seems to be mainly due to the lower affinity of this antibody for the FcγRIIa alloforms. The mAb MA73 (2ZC115), which is negative on intact cells (Table 1), specifically reacts with both FcγRIIb isoforms in cell extracts. Using a series of FcγRIIb2 mutants [2] lacking various numbers of amino acids from the carboxy-terminal end of the receptor we found that this mAb does not recognize FcγRIIb2 mutants lacking nine amino acids at the carboxy terminal end. In contrast, changing the Tyr273-residue into a Phe residue

does not influence the reactivity of mAb MA73 (2ZC115).

References

1. Budde, P., Bewander, N., Weinrich, V., and Frey, J. *Eur. J. Cell Biol.* 64, 45 (1994).
2. Engelhardt, W., Gorczyca, H., Butterweck, A., Mönkemann, H., and Frey, J. *Eur. J. Immunol.* 21, 2227 (1991).
3. Greenman, J., Tutt, A. L., George, A. J. T., Pulford, K. A. F., Stevenson, G. T., and Glennie, M. J. *Mol. Immunol.* 28, 1243 (1991).

M8.3 Binding heterogeneity within the CD32 panel of mAb

INGRID E. VAN DEN HERIK-ODDIJK, NOMDO A. C. WESTERDAAL, MASJA DE HAAS, LAURA KEMPER, PETER J. A. CAPEL, ALBERT E. G. KR. VON DEM BORNE, and JAN G. J. VAN DE WINKEL

The CD32 molecule (FcγRII) represents a 40-kDa low-affinity receptor for IgG, and is encoded by three genes, FcγRIIA, IIB, and IIC, all localized on chromosome 1q23-24. The transcripts derived from genes IIA (FcγRIIa) and IIB (FcγRIIb) differ both in their signal peptides and cytoplasmic domain-encoding regions, whereas extracellular and transmembrane encoding regions are ~92 per cent homologous [1]. The FcγRIIC gene has been characterized as a result of a cross-over event between the IIA and IIB genes [2], and the transcript of gene IIC was found to be identical in its signal peptide and extracellular and transmembrane-encoding regions to FcγRIIb, whereas the cytoplasmic region-encoding domain was identical to that of FcγRIIa [3]. FcγRIIA is, furthermore, polymorphic and two allotypes have been defined that differ by a single amino acid (aa) at position 131 within the second Ig-like domain, where an arginine or histidine is found. These two allotypes differ in their ability to bind mouse (m) IgG1 complexes: FcγRIIa-R131 interacts effectively with mIgG1 (previous name: high-responder FcγRIIa), in contrast to FcγRIIa-H131 (previously: low-responder) [4]. Expression of the FcγRIIA gene is found on monocytes, neutrophils, and platelets [5], and evidence has been presented for (low-level) expression on B lymphocytes [6]. Additional diversity is found within the FcγRIIB subclass, which comprises three isoforms: FcγRIIb1,

IIB2, and IIB3. The FcγRIIb2 isoform is identical to IIB1 except for the lack of a 19-aa insert in the cytoplasmic region (due to alternative splicing of the C1 exon). FcγRIIb3 is almost identical to IIB2, but lacks information for the putative signalase cleavage site, due to an alternatively spliced S2 exon [3]. The FcγRIIb1 and IIB2 transcripts were both found to be expressed on B lymphocytes [6].

In order to assess the reactivity of the CD32 Workshop panel monoclonal antibodies (mAb) with the different FcγRII molecules, we generated a panel of stable transfectants. Three different mouse cell lines were used for transfection, in order to avoid reactivity with endogenous human FcγRII molecules on cells, and to minimize other cell type-specific effects (for example, variation in glycosylation patterns). 3T6 fibroblasts were transfected with cDNAs encoding FcγRIIa-H131, IIA-R131, and FcγRIIb1* [4,7]. Of these, FcγRIIb1* is identical to FcγRIIb1 except for one aa difference at position 11 within the cytoplasmic tail, where a tyrosine (IIB1) is replaced by an aspartic acid (IIB1*). Moreover, these cDNAs, as well as FcγRIIb1 and IIB2 cDNAs, were expressed in a mFcγR-negative and surface IgG2a-positive mouse B-cell line IIA1.6 [6]. The third cell type we used for transfection was mouse T-cell line RMA-S, which binds anti-mouse FcγRII/III mAb 2.4G2 [8] and, thus, expresses endogenous mouse FcγR. These

BEST AVAILABLE COPY